

Common Name: Banai-banai

Scientific Name: Radermachera quadripinnata (Blanco) Seem.

Family: Bignoniaceae

Distribution: Borneo, Malay Peninsula, Moluccas, Nicobar Isls, Philippines, Sulawesi,

Sumatra, Thailand. BILIRAN, BOHOL, CAMIGUIN, CEBU, CULION, HOMONHON, LEYTE, LUZON: Albay, Bataan, Benguet, Cagayan,

Camarines Sur, Cavite, Ilocos Norte, Ilocos Sur, Isabela, Laguna, Mountain

Province, NCR, Pangasinan, Quezon, Rizal, Zambales, MINDANAO: Agusan del Norte, Lanao, Lanao del Sur, Misamis, Zamboanga,

Zamboanga Sibugay, MINDORO, NEGROS, PALAWAN, POLILLO, SAMAR

DAO/IUCN category: Least Concern



Description: Banai-banai is an evergreen tree growing up to 40 m tall. It produces clusters of pink-purplish-orange flowers and fruit pods that are elongated capsules which splits, shedding numerous tiny, silky, winged seeds, when ripe. This tree used in traditional medicine to treat a variety of ailments is recently gaining demand as an ornamental tree in parks and gardens.

Source: SUTCLIFFE, L.G. and MALABRIGO JR., P.L. 2020. A Guide to Westgrove's Native Trees. Vol. 1. Silang, Cavite: Ayala

Westgrove Heights Homeowners Associations, Inc. 30 p. https://www.philippineplants.org/Families/Bignoniaceae.html

 $Image\ Source: \ http://herbarium.bh.cornell.edu/CUBIC_IMAGES/phytoimages/pelserpb/4_7_17/7apr17/DSC_1355.jpg$